

J. Jason Wendel, M.D.
Dr. J. J. Wendel Plastic Surgery
2103 Crestmoor Road
Nashville, TN 37215
www.drjjwendel.com

Postoperative Instructions

LOWER FACE LIFT – NECK LIFT – MINI FACE LIFT

This is an outpatient surgery.

Have someone drive you home after surgery and help you at home for 1-2 days.

Get plenty of rest; follow balanced diet.

Decreased activity and pain medication may promote constipation, so you may want to add more fresh fruit to your diet, and be sure to increase fluid intake.

Take pain medication as prescribed. Do not take aspirin or any products containing aspirin until approved by your surgeon.

Do not drink alcohol when taking pain medications.

If you are taking vitamins with iron, resume these as tolerated.

Do not smoke, as smoking delays healing and increases the risk of complications.

Activities

Walk as soon as possible, this helps reduce swelling and lowers the chance of blood clots.

Do not drive until you are no longer taking narcotic pain medications.

Plan on taking it easy for the 5-7 days.

No strenuous activities, including sex and heavy housework, for 2 weeks.

Walking and mild stretching are okay.

No swimming for 4 weeks.

Return to work in about 1-2 weeks.

Incision Care

If a drainage tube is placed, you may shower 48 hours after removal.

You may bathe 48 hours after surgery. Avoid steam baths and saunas for 4 weeks.

Avoid exposing scars to sun for at least 12 months.

Always use a strong sunblock, if sun exposure is unavoidable (SPF 30 or greater).

Keep steri-strips in place.

Keep incisions clean, dry and inspect daily for signs of infection.

Keep head elevated; sleep with head on 2 pillows or sleep in a recliner.

Do not use hair dye or permanent hair solutions for two months.

You may use cold compresses for comfort and to help decrease the swelling.

Hair is generally shampooed on the 2nd or 3rd postoperative day. The area of sutures must be washed gently each time.

Keep hair dryer temperature set on low or no heat, as you temporarily may not have feeling in operated areas. You do not want to burn your skin.

You may use makeup after the sutures are removed; new facial makeup can be used to conceal bruising, but not on the incisions. Gently remove all

makeup prior to applying new makeup.

What To Expect

Maximum discomfort should occur in the first few days, improving each day thereafter.

Bruising, swelling, numbness, and tightness of skin for 1-2 weeks.

You will be sent home wearing a chinstrap. This provides compression and support.

You may experience tightness in the neck for 2-4 weeks.

Skin may feel dry and rough for several months.

Neck may look and feel strange or seem distorted from the swelling.

Appearance

Pale, bruised and puffy face or neck for several weeks.

Wearing scarves, turtlenecks and high-collared blouses masks the swelling and discoloration.

By the third week, you will look and feel much better.

Final result is not fully realized for approximately 3 months.

Follow-Up Care

Sutures are usually removed in 5-7 days.

If a drainage tube was inserted, it will be removed in 1-3 days.

Frequent follow-up visits should be planned.

When To Call

If you have increased swelling or bruising.

If swelling and redness persist after a few days.

If you have increased redness along the incision.

If you have severe or increased pain not relieved by medication.

If you have any side effects to medications; such as, rash, nausea, headache, vomiting.

If you have an oral temperature over 100.4 degrees.

If you have any yellowish or greenish drainage from the incisions or notice a foul odor.

If you have bleeding from the incisions that is difficult to control with light pressure.

If you have loss of feeling or motion.

For Medical Questions, Please Call:

(615) 921-2100 Monday-Thursday 9:00 AM-5:00 PM and Fridays 8:00 AM – 12:00 PM. After hours and on weekends, the answering service will contact Dr. Wendel or the plastic surgeon on call.