

J. Jason Wendel, M.D.

Dr. J. J. Wendel Plastic Surgery
2103 Crestmoor Road
Nashville, TN 37215
www.drjjwendel.com

Postoperative Instructions

EYELID LIFT SURGERY (BLEPHAROPLASTY)

This is an outpatient surgery. If you choose to have blepharoplasty done under local anesthesia, the procedure may be done in the office.

Have someone drive you home after surgery and help you at home for 1-2 days.

Get plenty of rest; follow balanced diet.

Decreased activity and pain medication may promote constipation, so you may want to add more fresh fruit to your diet, and be sure to increase fluid intake.

Take pain medication as prescribed. Do not take aspirin or any products containing aspirin unless approved by your surgeon.

Do not drink alcohol when taking pain medications.

If you are taking vitamins with iron, resume these as tolerated.

Do not smoke, as smoking delays healing and increases the risk of complications.

Sleep with your head elevated 45 degrees for several days to minimize swelling. Some patients find sleeping in a recliner helpful.

Activities

Walk as soon as possible, this helps reduce swelling and lowers the chance of blood clots.

Do not drive until you are no longer taking narcotic pain medications.

Limit activities to light duty for 3-5 days.

No lifting greater than 5 pounds for 1-2 weeks.

Avoid activities that raise your blood pressure, including bending, lifting and rigorous sports for 2-3 weeks.

No swimming for 2-3 weeks.

Return to work in 7-10 days.

Incision Care

Avoid exposing scars to sun for at least 12 months.

Always use a strong sunblock, if sun exposure is unavoidable (SPF 30 or greater).

Keep steri-strips in place.

Keep incisions clean, dry and inspect daily for signs of infection.

Use cold saline compresses for the first 24-48 hours to reduce swelling and bruising.

You can shampoo your hair 1-2 days after surgery.

Wait a minimum of 2 weeks before wearing contact lenses. Lubricate with lubricating eye drops frequently.

You will have sensitivity to sunlight, wind and other irritants for several weeks.

Wear sunglasses and eyelid cream that contains sunblock.

What To Expect

Maximum discomfort should occur in the first few days, improving each day thereafter.

Expect temporary swelling of the eyelids, tightness of lids, bruising, dryness, burning, and/or itching of eyes.

You may have oozing or crusting of eyes for approximately 1 week.

Use Natural Tears (OTC eye drops) or Lacrilube (OTC ophthalmic ointment) as part of your post op treatment.

For the first few weeks, you may experience tearing, sensitivity to light, and double or blurred vision.

You may have difficulty closing your eyes when you sleep. Use ointment at night to protect your eyes.

Appearance

Bruising and swelling of the eyelids may last 1-2 weeks.

Healing is a gradual process and your scars may remain slightly pink for 6 months or more.

Tiny whiteheads may appear after stitches are taken out; can be easily removed by surgeon.

Facial makeup can cover up bruising after the sutures are removed.

Follow-Up Care

Sutures are usually removed in 5-7 days.

When To Call

If you have increased swelling or bruising.

If swelling and redness persist after a few days.

If you have increased redness along the incision.

If you have severe or increased pain not relieved by medication.

If you have any side effects to medications; such as, rash, nausea, headache, vomiting.

If you have an oral temperature over 100.4 degrees.

If you have any yellowish or greenish drainage from the incisions or notice a foul odor.

If you have bleeding from the incisions that is difficult to control with light pressure.

If you have loss of feeling or motion.

For Medical Questions, Please Call:

(615) 921-2100 Monday-Thursday 9:00 AM-5:00 PM and Fridays 8:00 AM – 12:00 PM. After hours and on weekends, the answering service will contact Dr. Wendel or the plastic surgeon on call.