

J. Jason Wendel, M.D.
Dr. J. J. Wendel Plastic Surgery
2103 Crestmoor Road
Nashville, TN 37215
www.drjjwendel.com

Postoperative Instructions

PREVENTING POST SURGICAL CONSTIPATION

Preventing or alleviating constipation usually requires that the patient do the following:

- Increase fluid intake, dietary fiber and activity level.
- Establish and maintain regular bowel habits.

Below is a protocol to prevent and manage constipation caused from pain medication:

1. Prevention – Take 2 Senokot and 2 Colace tablets each night at bedtime.
This should produce a bowel movement the following morning.
(If no bowel movement in 2 or 3 days, follow instructions from item 2.)
2. Take 2 Senokot and 2 Colace tablets in the morning and also at bedtime. If you usually have a bowel movement daily, you can add Milk or Magnesia 30mL at bedtime.
(If no bowel movement in 2 or 3 days, follow instructions from item 3.)
3. Take 3 Senokot and 3 Colace tablets in the morning and also at bedtime.
(If no bowel movement in 2 or 3 days, follow instructions from item 4.)
4. Take 4 Senokot and 4 Colace tablets in the morning, at midday and also at bedtime.

If you have not had a bowel movement after 4 total days on this protocol, call the office at (615) 921-2100 and speak to a nurse.

For Medical Questions, Please Call:

(615) 921-2100 Monday-Thursday 9:00 AM-5:00 PM and Fridays 8:00 AM – 12:00 PM. After hours and on weekends, the answering service will contact Dr. Wendel or the plastic surgeon on call.